

Body condition scoring

The reference guide for New Zealand dairy farmers





For more information visit dairynz.co.nz

DairyNZ Corner Ruakura and
Morrinsville Roads Private Bag 3221
Hamilton 3240

Version 3 - 08/2024 DNZ10-102

Disclaimer

DairyNZ Limited ("DairyNZ", "we", "our") endeavours to ensure that the information in this publication is accurate and current. However, we do not accept liability for any error or omission.

The information that appears in this publication is intended to provide the best possible dairy farm management practices, systems and advice that DairyNZ has access to. However, the information is provided as general guidance only and is not intended as a substitute for specific advice. Practices, systems and advice may vary depending on the circumstances applicable to your situation. The information may also be subject to change at any time without notice. DairyNZ takes no responsibility whatsoever for the currency and/or accuracy of this information, its completeness or fitness for purpose.

©DairyNZ Limited 2024

Contents

Preface	4
Introduction	4
<hr/>	
Section 1 – Assessing BCS	5
Assessing BCS using the DairyNZ method	5
Why use the DairyNZ BCS method?	5
BCS versus liveweight	7
Breed variation and BCS	8
How to BCS	9
Who should score cows	13
When to score	14
How many cows to score	14
Tools to record scores	15
Locations to score	16
Animal welfare and BCS	18
When to take action	19
<hr/>	
Section 2 – BCS targets	20
BCS targets at calving and mating	20
Why use different targets for younger cows?	22
Why achieving BCS targets makes the difference	22
<hr/>	
Section 3 – Strategies to achieve targets	24
How to achieve BCS targets	24
• Increase feed allocation to lactating cows	24
• Reduce milking frequency in mid-late lactation	24
• Dry off cows early	25
• Feed dry cows for BCS gain	26
Other considerations	29
<hr/>	
Section 4 - The science of BCS	31
BCS targets - a balancing act	31
Impact of BCS	31
Factors that affect BCS	37
Summary	42
<hr/>	
Section 5 - Appendices	43
References	43

Preface

This resource is a companion to the DairyNZ Body Condition Scoring Made Easy field guide. It aims to provide farmers and rural professionals with advice on body condition score (BCS) targets, the effect of management strategies on BCS, and an overview of how to use the DairyNZ BCS method to get accurate and consistent results whenever cows are scored.

This reference guide summarises a scientific review on the effect of BCS on key productivity-related traits and animal welfare, and the relationship between farm management, cow genetics and nutrition on BCS change. The review was compiled by an international team led by DairyNZ scientists. It has been published in the Journal of Dairy Science (J. Dairy Sci. 92:5769–5801) and can be accessed, in full, at journalofdairyscience.org

Introduction

A cow's body condition score (BCS) is a reasonably accurate estimate of her energy reserves, and BCS targets at key stages of lactation have been identified to optimise dairy production systems. These targets are based on research studies undertaken in New Zealand and internationally over the last 40 years. These studies were summarised in a scientific review published in the Journal of Dairy Science led by DairyNZ scientists. The contents of this review confirmed and quantified the recognised effects of BCS on dry matter intake, milk production, reproduction, and cow health and welfare. In addition, there was evidence that BCS in early lactation can affect the sex of future calves and the productive and reproductive capacity of heifers yet to be born.

Irrespective of the system of farming (low or high input), cow genetics, or the country where the research was undertaken, productivity was optimised when mature cows calved between BCS 5.0 and 6.0 and when first and second calvers were 0.5 BCS units fatter than mature cows. That said, the productivity benefit declined with increasing BCS and the risk of metabolic health disorders at calving was greater, such that increasing BCS beyond 5.0 in mature cows and 5.5 in first and second calvers isn't justified. The recommended calving BCS targets provide a compromise between optimising energy reserves for milk production, while minimising cow health issues caused by over-fatness. To ensure calving BCS targets are achieved, the most appropriate management strategies are explored in this resource. Although target BCS recommendations tend to be herd-focused, the review also highlighted the need to consider individual cow BCS. Animal welfare is concerned with the state of the individual cow, not the average of the whole herd. Achieving the desired herd average target may still leave a proportion of cows that are too thin, or too fat. The Dairy Cattle Code of Welfare 2019 (New Zealand Ministry for Primary Industries) outlines the minimum standard for body condition score for dairy cows.

Farm management and nutrition in very early lactation (first four to five weeks after calving) has little impact on the speed at which a cow loses BCS or the amount of BCS lost. However, mid and late lactation management and nutrition, and nutrition when the cow is dry, influence how fast a cow gains (or loses) BCS. Management strategies that help to achieve BCS targets are presented in this resource. This resource should be used as a companion to the DairyNZ Body Condition Scoring Made Easy field guide.

Find the DairyNZ Body Condition Scoring Made Easy Field Guide at
dairynz.co.nz/bcs-field-guide



If you would like advice on your body condition scoring, contact DairyNZ at
dairynz.co.nz/about-us/contact-us

Assessing BCS using the DairyNZ method

- Body condition score (BCS) is a subjective assessment of a cow's energy reserves, rating the "body fat" of a cow on a scale of 1 to 10, where 1 is emaciated and 10 is obese.
- The DairyNZ BCS method helps standardise how cows are scored.
- When learning to body condition score it is important to calibrate the eye by scoring cows using the 'hands-on' method (see **page 9**).
- To ensure independent, accurate and consistent body condition scoring, use a Certified BCS Assessor.
- For most people, visually scoring more accurately than 1.0 BCS units requires formal training. Assessors can confidently score cows to 0.5 BCS units visually. Very experienced assessors can score to 0.25 BCS units by feeling the main body parts, but not by sight alone.
- For much of the year, an average BCS (and range) of the herd will be appropriate. At least 70 randomly selected cows should be scored in each herd. However, there are key times during the year where scoring all cows is important.
- The most important time to body condition score individual cows is during summer and autumn. Scoring all cows at this time will allow individuals to be managed differently (see **pages 20-23**) to ensure BCS targets at calving are met.

Why use the DairyNZ BCS method?

There are several different methods for assessing BCS with varying degrees of accuracy and consistency. The method developed and supported by DairyNZ is regarded as the standard (**Table 1**).

Introduced and developed over 40 years ago, this method is internationally recognised. It forms the basis of milk production, reproduction and animal health publications quantifying the effect of BCS on productivity and sustainability for New Zealand dairy systems.

The method is recognised and recommended by the Ministry for Primary Industries and embedded as Schedule II in the Dairy Cattle Code of Welfare 2019.

Dairy cattle in New Zealand vary considerably in size and frame compared with dairy cows in other parts of the world. There is also variation in the way New Zealand dairy cows store fat reserves across their bodies. The DairyNZ method takes this variation into account by assessing fat reserves at multiple points across the frame of the animal to correctly assess overall BCS.

Once mastered, this method is fast and accurate and is supported by additional BCS resources.

Explore DairyNZ BCS resources and tools at dairynz.co.nz/animal/body-condition-scoring



Important body points to consider when condition scoring cows




Table 1. Quick guide to body condition scoring

Body part	Descriptor
Backbone	Is it flat or is there a ridge? Can you see or easily feel notches?
Long ribs	Can you see or easily feel the ribs? If visible, how many can you see?
Short ribs	Can you see the short ribs? What do they feel like? Are the rib ends sharp or rounded?
Hip bones	Are the hip bones rounded or angular?
Rump	Is the area between the pins and hip bones, flat, sunken or hollow?
Pin bones	Are they pointed, tap-like or rounded?
Tailhead	Is there a hollow between the tail head and pin bones? Is it a deep 'V' or shallow 'U' shape?
Thigh	Is the area indented, flat or rounded? Is the muscle structure defined?

BCS versus liveweight

Liveweight (Lwt) change is not a good indicator of BCS change. This is why BCS systems were developed globally. Factors such as breed differences, cow-to-cow variation in size within a breed, change in gut fill, pregnancy, and heifer and second calver growth prevent the use of Lwt as an accurate measure of an individual cow's energy reserves. New Zealand research results indicate that Lwt change only explains 25% of the variation in BCS change.

Despite its limitations, many farmers measure Lwt as an indication of how much BCS a cow has gained or lost. As a guide, the average amount of weight gain/loss for every unit of BCS change is equivalent to 6.58% of a cow's total Lwt. The BCS system is less relevant to youngstock, which should be regularly weighed to monitor liveweight gain (**Table 2**).

Table 2. Average Lwt change per BCS unit for common breeds are as follows:

Jersey (425 kg Lwt)	1 unit change in BCS = 28 kg
Crossbred (475 kg Lwt)	1 unit change in BCS = 31 kg
NZ Holstein-Friesian (500 kg Lwt)	1 unit change in BCS = 33 kg
Overseas Holstein-Friesian (550 kg Lwt)	1 unit change in BCS = 36 kg

Even with training and experience, an assessor can only score cows to a 0.5 BCS unit visually with confidence. Remember, 0.5 BCS units is only 14-18 kg Lwt. Herd average BCS can be reported to 0.1 BCS units.



Breed variation and BCS

Although there are significant differences in size and angularity between the common New Zealand dairy breeds, the same guidelines for BCS assessment can be applied to all breeds.

Jerseys are typically smaller framed, with a comparatively narrow body and prominent hip bones.



Crossbreds and New Zealand-type Holstein-Friesians are medium framed animals that carry fat reserves evenly over the body.



New Zealand-type Holstein-Friesians are generally shorter, rounder and blockier than the larger overseas Holstein-Friesian type.



Overseas Holstein-Friesian are more angular, appear thinner and tend to carry more body fat over the long ribs and less around the tailhead than New Zealand-type Holstein-Friesians.



How to BCS


Body condition scoring is a simple process. However, it is important to calibrate the eye by first condition scoring cows 'hands on'. To score, line up 15 cows with a range of BCS in a vet/AB race.

- Put your hands on the eight body parts of the cow outlined below. Feel the amount of fat cover over the various body points and the difference between the cows.
- Note that gut fill can give a visual impression of condition over the ribs, therefore it is important to feel the amount of fat cover.
- The overall BCS of the cow will be the average of the scores for the eight body parts.

The eight body parts to score:


Backbone

- When assessing the backbone, look at the ridge formed by the spine above the short ribs (loin).
- From the rear of the animal, is the backbone ridged and prominent or rounded and flat?
- From the side of the animal, are notches formed by the individual vertebrae visible or is the ridge of the back smooth?
- If no notches are visible, then BCS for the backbone is greater than 4.0. If notches are visible, then BCS is 4.0 or less.




Long ribs

- As BCS falls, long ribs gradually become more apparent. The last rib closest to the tail is the first to be seen and then more ribs appear toward the head as BCS falls (this area in particular can be influenced by shadow, cow colour and whether the cow is wet or dry).
- Can the ribs be easily seen and, if so, how many? If the long ribs cannot be seen from a distance can they be felt?
- If long ribs are not visible, the BCS for this part is more than 5.0. If long ribs are visible down to the level of the udder, then the BCS for this part is below 3.0. Note, if the overall BCS for the cow is below BCS 3.0, then urgent remedial action must be taken.




Short ribs

- The 'short ribs' of a cow are not actually ribs at all; they are part of the spine called transverse processes. As BCS falls, the ends of these bones become more pronounced and sharper to the touch. Short rib scores can be influenced by rumen fill. When the rumen is full, the "triangle" below the short ribs is filled on the animal's left-hand side, giving the impression that the animal is carrying more condition than it actually is.
- Can the short ribs be easily felt? Are the ends of the ribs sharp, smooth or rounded?
- When scoring cattle, it is important to view the right-hand side of the animal so as not to be influenced by rumen fill. If the short ribs have a "gloved fingers" appearance, BCS for this part is less than 4.0.




Hip bones

- Sometimes referred to as 'hooks' or 'hook bone'. Are the hip bones angular and sharp or are they smooth and rounded?
- Place a cupped hand over the hip bone. If the hip bone feels rounded, BCS for the hip bones will be 5.0 or more.
- If there are depressions either side of the hip bones, BCS will be 3.5 or less.




Pin bones

- The pin bones protrude either side of the tailhead and become rounded with fat deposits as BCS increases.
- Are there dimples/indents between the points of the pin bones? Are the pin bones angular or rounded?
- If the pin bone forms a tap-like shape, BCS for the pin bones is 4.0 or less.




Tailhead

- The tailhead is the area between the pin bones. As BCS increases, fat deposits change the shape and depth of the hollows between the pin bones and the tail itself.
- Does the indentation on either side of the tail base form a 'U' or a 'V' shape?
- Use your fingers to feel the degree of indentation. If the tail head feels like a shallow 'U' shape, BCS for the tailhead is 4.0 or above. If the tail head feels like a deep 'V' shape, BCS for the tailhead is 3.0 or below.




Rump

- The rump is the area between the hip bone and the pin bones. As BCS increases, this area fills from hollow to concave, then flattened to bulging and 'beef' like.
- Is the area deeply sunken, hollow, or flat?
- Draw an imaginary line from hip to pin bones following the contour of the rump. Is the line flat, dished, or does it form an open 'V'? If the rump forms an open 'V', BCS for the rump is 3.0 or less.



Thighs

- Best viewed from the side and rear of the animal, the thigh can be the hardest area to assess.
- As BCS increases, the areas below the rump and above the hocks fill out. Are they indented, flat or rounded?
- If the muscle structure of the thigh is visible, BCS is less than 4.0.



Who should score cows

Farmers and rural professionals can score cows. But while body condition scoring is a simple process, it requires training, practice and regular calibration to ensure accuracy and consistency. Most people need formal training to visually score cows with greater accuracy than 1.0 BCS units. With training, assessors can confidently score cows to 0.5 BCS units visually.

Accurate and consistent BCS scores can then be used to support good management decisions, providing cows with the best chance to meet BCS targets and optimise production, health and welfare outcomes. Scoring bias can also occur when you are scoring cows that you see every day. An independent scorer can ensure this is avoided.

The best way to ensure independent, accurate and consistent body condition scoring using the DairyNZ method is to use a Certified BCS Assessor. Potential BCS assessors attend a training workshop and then complete an assessment. If successful, assessors are certified for 12 months. To remain certified, assessors must attend a formal calibration event every 12 months.

What if I'm not a certified BCS assessor?

It is essential for the scorer to ensure that they are consistent each time they score. Failure to regularly calibrate, leads to variation in scoring, which can lead to poor management decisions.

For those not certified, there are options to improve scoring accuracy:

- Use the DairyNZ Body Condition Scoring Made Easy field guide as a reference to confirm the score given as well as other online resources from DairyNZ.
- Score 10 to 15 animals 'hands-on' in vet/AB race before herd scoring. 'Hands-on' involves assessing each of the eight body parts of the cow.
- Score cows alongside your certified vet/consultant and compare results.
- Attend a DairyNZ BCS training workshop.
- Consider becoming a Certified BCS Assessor.

Visit DairyNZ BCS training events at dairynz.co.nz/support/training/bcs-training-events



When to score

Summer and Autumn: Are the most important times to body condition score cows. This allows individual groups of cows to be managed differently to ensure BCS targets at calving are met.

Pre-Christmas assessment: Determines if cows have gained BCS since the planned start of mating and allows a management plan for the autumn to be set up.

Mid-February to mid-March: Assessing cows determines if the management plan is working or if some cows should be dried off early, milked once-a-day (OAD), or preferentially fed.

Pre-calving: Assessing BCS (in springer mob) determines if the autumn management plan achieved the calving BCS targets and whether there is a need to re-assess management practices for next autumn.

Planned start of mating: Helps determine how much BCS cows lost between calving and mating. This will identify if getting cows back in calf is at risk as well and whether anything should have been done differently to prevent the BCS loss.

How many cows to score

The number of cows you need to score will depend on what you want to achieve. For much of the year an average BCS (and range) of the herd may be appropriate. The greater the percentage of animals assessed, the more accurate the result. For groups greater than 100 cows, a minimum of 70 cows should be scored when determining a herd average. These cows should be selected at random. With practice, scoring 70 cows will take less than 30 minutes. Scoring all cows in the herd allows individual management and is most important during summer and autumn.

Tools to record scores

BCS app

The BCS app allows you to score cows on your smartphone and have ready access to the results. Use the app to:

- Calculate the average score for the herd, and store and review over time.
- Score multiple herds and create a report.
- Flag an individual cow and record her tag for later reference.

To download the app, visit dairynz.co.nz/bcs-tracker



DairyNZ BCS recording sheet

If you prefer a paper method, the DairyNZ BCS recording sheet can also be used (Figure 1).

Blank recording sheets can be downloaded from dairynz.co.nz/bcs-chart



Figure 1 - BCS recording sheet

This herd size =		126 cows						Date:		04-06-2024	
Condition score	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	7.0	
Number of cows		13	27	33	27	14	8	4			
Total BCS		39	94.5	132	121.5	70	44	24			
% of sample (Number of cows in BCS group ÷ Total Cows (A))		10.3%	21.4%	26.2%	21.4%	11.1%	6.4%	3.2%			
									Totals		
									126	A	
									525	B	
Average Condition Score:		Total CS		Divide by		Total Number of cows		=	Average BCS		
		B 525		÷		A 126		=	4.2		

Locations to score

Cows can be scored in different locations but there are pros and cons for each location. Body condition scoring should be done from the rear right side to minimise the effect of rumen fill, which can reduce the visibility of long and short ribs on the left-hand side of the cow when her rumen is full.

In a vet race

A vet race is the best location to provide an accurate BCS score, as you can get your hands on the eight body parts of the cow, ideal for calibrating. When scoring in a race, keep in mind the following points to maintain scoring accuracy:

- Cows need to be in single file, with the right side of the cow being unobscured (no distracting shadows, fences, gates, or other stock).
- Cows should be stationary, or moving very slowly, whilst being inspected for scoring.

This method is time consuming. It is good for learning and calibrating but not for scoring large numbers of cows.

In the paddock

Scoring in the paddock allows for more cows to be scored in a shorter period of time, and is a good option once you are more confident with the hands-on method.

When scoring, adapt to the flight distance of the herd being scored. If you are more than seven metres away from the cow being scored your accuracy is at risk. If the average flight distance of the mob/herd is greater than seven metres, then the BCS assessment should be conducted in another location.

Take care not to score the same cow more than once; entering cow ID in the BCS app can help avoid this. It is difficult to score the whole herd in the paddock.

On the feedpad

Walking quietly behind cows as they feed on the pad can also be a useful location to BCS cows. This location may allow for scoring a large number of cows, provided they are settled and there is adequate space for the assessor to move without disturbing them.


In the cowshed

Scoring in the cowshed can be a practical and convenient way to score large numbers of cows, however it is important to score from a position where you are at the same level or slightly above the cow being scored (e.g. while standing on a raised platform).

Scoring cows while they are in the bail can reduce the accuracy of scoring. In most situations the pit/bail configuration and presence of the rump and kick rail makes it difficult to see and assess all eight body parts required for body condition scoring. Inadequate lighting can also have a substantial effect on the ability to score accurately.

Rotaries

The limited view from the pit makes accurate BCS assessment difficult. Scoring from a rotary milking platform is acceptable, provided the scorer is on a raised platform. From this position the scorer can view the pin bones, hip bones and tailhead and can make a reasonably accurate assessment of the backbone, short ribs and rump. The thighs and long ribs will be obscured to some degree by the rails. The rotary speed needs to be sufficiently slow to enable the scorer enough time to score each cow. If necessary the rotary should be stopped periodically to allow scoring as needed. It may be possible to score all cows in the herd in this location.



Herringbones

Scoring cows in a herringbone dairy without a mobile raised platform/trolley should be avoided, due to reduced accuracy. However, cows of concern can be identified by standing at the cows' head and looking over the backline and making an assessment based on the backline, hip bones, short ribs and top of the long ribs. Once drafted into a race/yard, cows can then be accurately scored. If an average herd BCS is required, a random selection of cows can be drafted out into a race/yard and scored there. In general, it is not practical to score the whole herd in this location.



Animal welfare and BCS

As the person in charge of looking after animals, it is your legal obligation to ensure that any animal under your care meets the minimum standards outlined in the relevant codes of welfare, published by the New Zealand Ministry for Primary Industries. When the body condition score of any animal falls below 3 (on a scale of 1-10), urgent remedial action must be taken. Additionally, some NZ milk companies require farmers to take action sooner than this, and cease supplying milk from cows that are below BCS3.5 until they have regained condition.


For information on your obligation with regard to BCS see [Code of Welfare: Dairy Cattle 2019](#) (pg 44-46).

Urgent remedial action

Urgent remedial action may include providing additional feed, seeking veterinary advice, or on-farm slaughter. Note that animals below a BCS of 3 cannot be transported to a processor or saleyards. They can only be transported to an alternative property for additional feed with a current conditional transport certificate from a veterinarian. The flow chart below provides a guide about transport restrictions; it does not guarantee acceptance of livestock for transport or processing. For further information on your obligations for pre-transport selection, refer to dairynz.co.nz/animal/animal-welfare/transporting-stock

Transport restrictions based on BCS

Decision making tree



When to take action

Example of BCS 2.5



Table 3. BCS 2.5 – what to look for

Backbone	Very prominent, with visible notches that can easily be counted.	
Long ribs	The long ribs may be visible as far down as the top of the udder.	
Short Ribs	Clearly visible and easily counted. The short ribs will have very sharp edges.	
Hip	Deep depressions on the side of the hip bone.	
Pins	Very clear, three pronged tap formation.	
Tailhead	Very prominent and angular, with a very deep 'V' shape.	
Rump	Severely depressed (hollow).	
Thigh	No visible fat and very little muscle.	



BCS targets at calving and mating

At calving

- Mature cows should calve at a BCS of 5.0.
- First and second calvers should calve at a BCS of 5.5.
- With no more than 15% of the animals greater than, and not more than 15% of the animals less than, these targets.
- Note that first calvers (heifers) need to be at BCS 5.5 at 22 months of age.
- This requires monitoring the growth and health of youngstock, and a strategy to achieve targets, from weaning.

At mating

- Mature cows should be a minimum of BCS 4.0 at planned start of mating (PSM).
- First and second calvers should be a minimum of BCS 4.5 at PSM.
- With no more than 15% of the animals less than these targets.
- The average decrease in BCS for the whole herd between calving and mating should not be more than 1.0 BCS unit.
- Note that all animals should be gaining BCS before PSM.



Target for calving

Example of mature cow at BCS 5.0



Example of heifer at BCS 5.5



Minimum at planned start of mating

Example of mature cow at BCS 4.0



Why use different targets for younger cows?


BCS targets for first and second calvers are set at 0.5 BCS units higher than for mature cows. This is because younger cows:

- Tend to lose more BCS after calving as they are still growing to their mature weight and partition energy towards growth as well as lactation. Greater BCS at calving enables them to reach target BCS at mating.
- Tend to have a longer anoestrus period after calving than older cows, which leads to later cycling and lower conception rates. Aiming for a higher body condition at calving helps to reduce this risk.
- Are at a lower risk of metabolic disorders than older cows, so can tolerate calving at a higher body condition.

Why achieving BCS targets makes the difference

The BCS targets at calving represent the optimum 'sweet spot', and have a large impact on dry matter intake, milk production, reproduction, and health post-calving. There are as many negative consequences of being below target BCS at calving as there are for being above target (**Figure 2**).

Figure 2. Summary of the consequences of cows being below or above target BCS at calving



BCS at calving has the greatest influence on reproduction. Once cows have calved, it is very difficult to alter total, or rate of, BCS loss. Increasing feed allocation (pasture or supplements) does not impact on BCS loss until after 4-5 weeks of lactation. The main determinant of BCS loss after calving is BCS at calving, which is primarily influenced by feed and cow management in late lactation and during the dry and transition periods.

Decades of research in New Zealand and overseas has been used to define the body condition score targets at calving, which are:

- Mature cows (four years and older) should be at BCS 5.0 at calving.
- Two and three year-old (first and second calvers) should be at BCS 5.5.

These ensure that dry matter intake, milk production and reproduction are optimised, while minimising the risk of health and welfare issues.

What are the consequences of calving cows at BCS 4.5?

Cows calving at BCS 4.5 will produce less milksolids (~\$34 milk revenue at \$8.00/kg MS) and cycle three to four days later than they would have, had they calved at BCS 5.0. In addition, cows that calve at BCS 4.5 are more likely to be less than BCS 4.0 at mating and, therefore, have lower submission and conception rates.

There will always be a percentage of the cows that are not at BCS 5.0 at calving. The target is to have greater than 85% of the mature cows at BCS 5.0.


BCS loss after calving

Cows lose BCS naturally after calving. The fatter they are at calving, the more they lose. On average, a cow calving at BCS 5.0 will lose approximately 1.0 BCS unit after calving, while a cow calving at BCS 6.0 will lose approximately 1.5 BCS units and a cow calving at BCS 4.0 will lose 0.75 BCS units. This post-calving loss is primarily regulated by genetics and, provided the cow is healthy, cannot be influenced by management or nutrition during the first five weeks after calving. At this point, supplements will slow the rate of BCS loss and increase the rate of BCS gain. However, the effect of feeding on BCS gain is small while the cow is milking because she has been selected to turn feed into milk and not into BCS.



How to achieve BCS targets

Four main strategies are used to help achieve BCS targets at calving. Choice of strategy will depend on the gains required for individual cows, and practicalities of managing a specific farm. These are:



Start managing BCS gain as early as possible, as once into late lactation, it becomes more difficult to significantly alter BCS.

Drafting cows into mobs, based on BCS, age, and expected calving date allows the right mix of approaches to be applied to cows with similar gains in BCS required.

1. Increase feed allocation to lactating cows

Cows tend to preferentially partition energy to milk production at the expense of BCS during lactation. They also have higher energy demands compared to a dry cow, due to:

- 10-20% more energy required for maintenance, and
- 5-10% more energy required for walking and grazing activities.

Therefore, even in late lactation, providing additional feed to lactating cows will reduce/halt BCS loss, but is unlikely to cause significant BCS gain. Condition can be gained but will require more kg DM feed eaten than for a dry cow. If your pasture growth rates and covers allow, then increasing pasture allocation can be a cost-effective way of providing additional feed. Alternatively, supplementary feed can be used. The key focus when using supplements at this time of year should be on quality and cost, as the composition of supplement does not appear to affect BCS gain.

2. Reduce milking frequency in mid-late lactation

Once-a-day

Once-a-day (OAD) milking in mid-late lactation can be used to improve BCS at dry-off while achieving longer lactations. The strategy spares energy from activity (walking) and milk production and alters physiological processes towards BCS gain.

Previous OAD research in the Waikato has shown a 10-15% drop in daily milk production for herds that milked OAD for 2-3 months in mid-late lactation, compared with herds on twice a day (TAD) milking. As 60-70% of the season's production had occurred by this point, the impact on the whole season production is minimal (4-5%).

Milking cows OAD during this period improved energy status and BCS gain¹. But there was only a minimal decline in dry matter intake during the period of OAD milking, and a small reduction in feed required during the dry period, to achieve target BCS. If OAD milking starts in early summer, with reasonably well-fed cows, the expected BCS gain is only 0.2 BCS units/month.

The key to successfully using this strategy is to maintain a similar feed allocation to when cows were on TAD so there is energy available for BCS gain. If BCS gain is adequate, you may be able to increase lactation length and achieve similar total milk production as if cows continued to be milked TAD.

For farms that have a set day for dry-off, regardless of conditions, the opportunity to extend lactation length to compensate for the lower production during OAD milking will be lost. This needs to be considered when planning for dry off.

Note that switching to OAD milking can increase the bulk milk somatic cell count (SCC) for a short period of time. Individual cow SCC can double for two to three days, so the bulk milk SCC must be able to accommodate this doubling, and not reach penalty levels. After a few days, the bulk milk SCC will drop back down to around 50,000 cells/ml above where it was previously. One solution is to remove high SCC cows from supply and dry them off early to reduce the risk of grading before switching to OAD.

For farms that can run multiple herds in their system, switching just one herd to OAD milking in mid to late lactation is a good option. Cows that have the greatest need to gain condition before calving, such as low BCS cows, early calvers, and rising three- year-olds, can be milked OAD in a separate mob. This leaves the rest of the herd, often including the high SCC cows, on twice a day milking. Cows milked OAD for three months in late lactation tend to have 0.25 BCS units more condition at dry-off than those milked twice-a-day, provided cows milked OAD are well-fed.

3 in 2 milking

Compared to OAD milking, milking 3 times in 2 days (3in2) can minimise the milk production losses associated with OAD milking while still reducing milking frequency. However, the additional energy available for BCS gain will be dependent on the farm situation.

A 2019 DairyNZ study² examined the effect of 3in2 milking on production at different stages of lactation, using milking intervals of 12-18-18 hours. Preliminary results indicate only minimal production losses when implementing 3in2 (~5% total production for the period 3in2 was used), relative to the control cows milked TAD. This suggests limited additional energy available for BCS gain.

Cows milked 3in2 for the whole lactation were 0.25 BCS greater in May at dry off, compared to TAD milked cows. However, for cows milked 3in2 for shorter periods i.e., for the last three months in late lactation, there was a more limited effect on BCS gain, with cows gaining less than 0.1 BCS compared to cows milked TAD².

For the latest information on milking frequency, see dairynz.co.nz/milking/milking-intervals

3. Dry-off cows to support BCS gain

The length of the dry period required for cows to reach BCS targets at calving will be determined by:

- The amount of BCS needed to be gained
- The amount of feed available
- The type of feed that will be offered during the dry period

The number of days dry, summarized in **Table 4**, provides a guide as to the length of dry period needed for cows of different BCS in late lactation to reach BCS targets by calving. These calculations account for the 30-40 days in the dry period where no net BCS gain occurs.

Table 4. Drying off time based on cow BCS and time to calving to achieve target calving BCS

Body condition score		Days cow needs to be dry before calving	
Mature cow	Rising three-year-old	Autumn pasture with limited supplements or crops	Autumn pasture and/or high-quality crops supplement fed above maintenance)
3.0	3.5	160	120
3.5	4.0	130	100
4.0	4.5	100	80
4.5	5.0	70	60
5.0	5.5	42*	42*

Note: Includes 10 days when cows are being dried off and not gaining BCS, and 30 days when cows do not gain BCS before calving.

*DairyNZ recommends a minimum dry period of 42 days for all cows irrespective of BCS.

For cows at risk of not achieving body BCS targets, drying off early involves trading off autumn milk production to reach BCS targets before the winter. However, the milk foregone in the autumn is approximately equal to the production gained the following spring, so there is usually no net change in milk revenue.

Financial gains from improved BCS come from improved reproduction and milk production. The figures for your herd can be calculated using the InCalf Gap Calculator. See InCalf Reproduction Gap calculator at dairynz.co.nz/tools/incalf-reproduction-gap-calculator

There are additional benefits for cows calving at target BCS that cannot be easily valued economically, such as resilience of the system in a difficult spring and reducing the number of thin cows on the farm.

The simplest way to operate this strategy is to use a staggered approach to dry-off, targeting lower condition and/or younger cows for early dry-off, based on the guide in **Table 4**. Dry off dates for cows at different BCS can be calculated using the BCS Dry-Off calculator.

See Body Condition Score (BCS) Dry-Off Calculator at dairynz.co.nz/bcs-calc



4. Feed dry cows for BCS gain

Typically, a dry cow uses energy less efficiently than a lactating cow for BCS gain, requiring 72 MJ ME to store 1 kg LWT, compared to 50 MJ ME for a lactating cow.

But a dry cow tends to gain more BCS per unit of feed eaten, compared to a lactating cow because she does not have the same energy demand of milk production and has lower activity and maintenance requirements compared to a lactating cow. So, more energy is available to be partitioned towards BCS gain.

For example, feeding a lactating cow 3kg DM concentrate per day for 100 days during mid or late lactation would increase her BCS by approximately 0.12 BCS units. In comparison, feeding 3kg DM concentrate per day to a dry pregnant cow, for 60 to 100 days pre-calving, would increase her BCS by 1.0 BCS units.

During the dry period, time available, and the type of feed offered are the two key drivers of whether a cow will reach BCS targets at calving. For early dry-off to be successful, cows must be allocated enough energy through the dry period to support maintenance, pregnancy, activity, cold stress (if applicable) and BCS gain. This will be a minimum of 9-11 kg DM/ day, depending on breed and feed utilisation.

Time requirements


The most common reason for not reaching BCS targets is not allowing enough time. Remember that:

- Cows rarely gain BCS in the first one to two weeks after drying off. Sometimes they even lose BCS during this period because the active immune response involved in the drying off process in the udder has a high energy demand, and feed is often restricted to cows during this period to reduce milk production.
- Dry cows gain very little BCS in the last month before calving (less than 0.1 BCS units). This is due to the large energy demands and inefficient use of nutrients by the growing calf.
- On average, cows will only gain 0.5 BCS per month during the dry period if fed a pasture-based diet, and 0.6 – 0.7 BCS units per month if fed a diet containing high levels of supplement or high energy crops.

Figures 3a and 3b show how the time available and the **type of feed** offered during the dry period influence the cow's ability to reach BCS targets ^{3,4}. Each scenario shows the amount of feed required each day to provide for maintenance and pregnancy while gaining one BCS unit over the dry period, for a 450 kg crossbred cow.

If a long-enough dry period is provided i.e., 120 days or 90 days, a dry cow can consume sufficient feed to meet her maintenance and pregnancy requirements and to gain 1.0 BCS unit (**Figure 3a**). But if only a short dry period i.e., 60 days, is provided, daily intake requirements are greater than the amount she can typically consume (blue zone). Supplementing with a higher DM feed such as palm kernel extract (PKE) (**Figure 3b**) can increase the amount of energy she can consume but still won't overcome a dry period that is too short.

Figures 3a and 3b. Daily feed intake required to meet maintenance and pregnancy requirements, as well as gain one BCS unit in the dry period for a 450 kg crossbred cow, for different lengths of the dry period. The blue zone represents actual daily feed intakes that a dry cow can consume ^{3,4}.



Type of Feed

Energy in autumn pasture is used less efficiently for gaining BCS compared to energy from other feeds such as pasture silage, maize silage and (PKE). Research results suggesting that fat (e.g. PKE) may be used more efficiently than starch (e.g. maize silage) for BCS gain in dry pregnant cows. Pasture and maize silage are used with 20% more efficiency, while PKE is used with 40% more efficiency than autumn pasture³.

The feed requirements (kg DM) for 1.0 BCS unit gain from different feeds are summarised in **Table 5**.

The reason why different feeds have different effects on BCS gain is currently unknown, but the results are based on feeding studies in New Zealand. Note that feeding PKE to dry cows does not affect the fat evaluation index, so can be used through the dry period with no milk quality restrictions.

Table 5. Approximate amounts (kg DM) of 'commonly used feeds' required for a 1.0 unit increase in BCS for dry cows^{3,5}.

These are average feed requirements above the needs for maintenance, activity and pregnancy.

Breed	Kg Lwt*	Kg Lwt/BCS	Autumn Pasture	Pasture Silage	Maize Silage	PKE
Typical energy content of feeds (MJ ME/Kg DM):			11.5	10.5	10.5	11
----- Total Kg DM of feed required -----						
Jersey	350	23	145	110	115	85
Jersey	400	26	165	130	130	100
Crossbred	450	30	185	145	145	110
Holstein Friesian	500	33	205	160	160	125
Holstein Friesian	500	36	225	180	180	135

*Live weights are for the cow only and exclude the weight of the foetus.

Note that:

- The differences between maize silage, PKE and pasture silage are not statistically significant, which means we cannot say with full confidence that the numerical difference is real, however this is the best information available for New Zealand farmers.
- The daily amount of feed required during the dry period will differ, with greater intakes required later in pregnancy.
- Realistically, cows do not gain BCS during the last month before calving because of the energy demands of foetal growth.



Other considerations

The practicalities of feeding and managing dry cows will affect the success of achieving BCS targets at calving.

Feed wastage

Wastage needs to be accounted for when determining the amount of feed to offer. The way that a supplement is fed, such as on a feed pad or along the fence line, affects wastage. Rules of thumb for different ways to offer supplements results in these utilisations:

- In-shed feeding facility = 95% utilisation
- Feed-pad = 90% utilisation
- Trailer in paddock = 85% utilisation
- Silage on ground = 80% utilisation (dry conditions) to 60% utilisation (wet conditions)

Mob management

To achieve BCS targets, it is important to manage cows according to their needs. For herds with a range of BCS, manage cows in different mobs so that some cows do not become over-conditioned, while others fail to reach target. Cows that are very light at dry off (BCS 3.5 or less) may require preferential treatment, such as:

- Are the cows in poor condition for reasons other than feeding e.g., lameness, facial eczema, mineral deficiency, parasites, infection, disease, etc? If in doubt get the animals checked by a veterinarian.
- Can their energy needs be further reduced, or access to allocated feed be made easier, by running them in a smaller mob to reduce competition?
- Can they be offered a high-quality feed that is more efficient for BCS gain e.g., PKE, which can be fed ad-lib to dry cows? (Refer to DairyNZ Facts and Figures, pages 65-80 for more information at dairynz.co.nz/facts-and-figures)
- If feeding pasture, can they be fed to more generous grazing residuals (i.e., greater than 4 cm or 8 clicks on the rising plate meter)?

Winter grazing

Winter grazing can be used to achieve BCS targets, but care must be taken to ensure:

- Cows have access to enough good quality pasture to gain desired BCS (**Table 5**).
- The quality of winter grazing is assessed. If the feed is low quality (kikuyu or browntop) or feed utilisation is poor, BCS gain will be less than predicted.
- Cows are monitored throughout the dry period and fed differentially, if needed, to ensure all cows reach targets.

Winter grazing does have other benefits, especially if the milking platform is wet, by reducing pugging damage and protecting future pasture growth.

Wintering on crops (e.g. kale, swedes, fodder beet)

Winter crops are generally considered an economical option for feeding cows to gain BCS in regions where pasture growth is insufficient to meet cow demand during winter (e.g., South Waikato, Central Plateau, Canterbury and Southland). Swedes, kale and fodder beet have been the primary crops used.

Studies in New Zealand⁶ reported no difference in the final BCS of cows between different crops, when crops were allocated on a MJ ME basis, and efficiency of utilisation was allowed for:

- Transition onto the crop
- Access to, and utilisation of the crop
- Required supplements (e.g., minerals, fibre)

Determining accurate dry matter yields and allocating crop correctly remain the key factors influencing successful wintering on crops.

For more details on best practice use of crops, see dairynz.co.nz/feed/crops

Transition cow feeding

In the last two to three weeks before calving, different strategies should be implemented for cows that are at target, and for those that are not at their correct BCS target.

For cows that are under target:

- Continue feeding to 100% of their requirements, to better support their immune system during calving and maximise their chances to gain condition up to calving.

For cows that are at, or above, target:

- Restrict their energy intake to 90% of their requirements, to minimise the risk of metabolic problems occurring during and after calving.

These approaches are summarised in **Table 6**.

Table 6. Metabolisable energy (ME) and approximate dry matter intake (DMI) requirements for last 2-3 weeks before calving

Cows less than BCS 5.0 100% of daily requirement			Cows at or greater than BCS 5.0 90% of daily requirement		
LWT	ME requirements	DMI requirements*	LWT	ME requirements	DMI requirements*
350	85	7.7	350	76	6.9
400	94	8.5	400	84	7.6
450	102	9.3	450	92	8.4
500	111	10.1	500	100	9.1
550	119	10.8	550	107	9.7

* ME = 11 MJ/kg DM

Note that the DMI figures in Table 6 are 'down the throat' (feed eaten) figures and allowances need to be made for wastage of pasture and supplements. For example, a 450 kg cow at BCS 4.5 would require 9.3 kg DM daily down the throat. If cows were grazing and pasture utilisation was 90%, they would need to be offered 10.3 kg DM daily.

BCS targets – a balancing act

The pre-calving BCS targets of 5.0 for mature cows and 5.5 for first and second calvers aim to:

- Optimise dry matter intake.
- Optimise milk production and reproduction.
- Minimise risk of health and welfare issues around calving and during lactation.

This section looks at each of these factors in isolation, but the interactions between them need to be considered as they occur on farm. It discusses the background of how the targets have been derived, and their relevance in supporting a successful system.

Impact of BCS

1. BCS and dry matter intake

Fatter cows graze for less time than thinner cows after calving. Therefore, the greater the BCS at calving, the lower the DMI and the greater the loss of BCS (negative energy balance) post-calving.¹

This is due to hormones acting on the brain that reduce the cow's drive to eat. As a result, fatter cows at calving are at greater risk of ketosis, fatty liver, and milk fever (see BCS and health, **page 35**).

As calving BCS establishes the profile of BCS change throughout lactation, it can have lactation-long implications for cow dry matter intake. Data from grazing cows in New Zealand suggests that for every 1 unit increase in BCS at calving, cows consume 0.5-1kg DM less per day after calving.

The negative effect of BCS on DMI is the same, whether the increased BCS is a result of cow genetics or management¹.

- If cows are naturally thin (genetics), they have a greater drive to eat.
- If cows are thin because of an earlier shortage of feed (management), they have a greater drive to eat.

Achieving target BCS at calving is a compromise between:

- Wanting cows with sufficient energy reserves for milk production and fertility outcomes
- While not being too fat, which reduces intake and increases risk of metabolic disease after calving.


2. BCS and milk production

The loss of BCS and mobilised body tissues in early lactation provides fatty acids for milk fat synthesis and amino acids for milk protein synthesis. Due to the greater loss in BCS in fatter cows, milk fat and protein yields increase with increasing calving BCS, up to BCS 6.5

The increase in milksolids production gets smaller as BCS at calving increases (**Figure 4**)

- An increase in calving BCS from 3.0 to 4.0 increases milksolids production by 17 kg/cow
- An increase in calving BCS from 4.0 to 5.0 increases milksolids production by 12 kg/cow
- An increase in calving BCS from 5.0 to 6.0 increases milksolids production by 6 kg/cow

Figure 4. Relationship between BCS at calving and annual milksolids production²



Data for these associations were estimated from 2,500 lactation records, collected from almost 900 cows², managed under tightly controlled grazing management systems. Associations were modelled between BCS at calving and BCS changes after calving with milk fat and protein production in the following lactation.

To optimise milk production, reproduction and health, cows should calve at a BCS of 5.0 (5.5 for two- and three- year-olds) and the BCS loss between calving and mating should be limited to a maximum of 1.0 BCS.

3. BCS and reproduction

Fertility is affected by:


- BCS at calving,
- BCS at planned start of mating (PSM),
- loss of BCS in early lactation, and
- whether cows are gaining liveweight before and during mating.

These relationships are based on analysis of 2500 whole-lactation records, collected from almost 900 cows³ managed under tightly controlled grazing management systems over a number of years. Associations were modelled between BCS at calving, BCS changes after calving, and various reproductive outcomes in the following lactation.

Thinner cows at calving cycle later

- A cow that calved at BCS 4.0 was less likely to have a pre-mating heat recorded than a cow calving at BCS 5.0 (**Figure 5**).

Figure 5. Relationship between calving BCS and the likelihood of cows cycling before planned start of mating (PSM)³.




Limitations of the dataset

- Only small numbers of cows were at a BCS below 3.5 or above 6.0 at calving. For this reason, the relationship for data within the 95% confidence interval is shown for **Figures 5, 6, and 7**.
- Less than 5% of the cows were at a BCS of 6.0 or over at calving, so the health and reproductive consequences for animals above target BCS are not well represented.
- Animals were removed from the herd due to health problems, potentially resulting from high or low BCS. This means that the full impact of BCS at the extremes on fertility and animal health may not be fully represented.

Cows that lose more BCS in early lactation have lower pregnancy rates

- A cow at BCS 3.0 at PSM was less likely to get in calf in the first six weeks of mating than a cow at BCS 4.0 at PSM (**Figure 6**).


Figure 6. Relationship between BCS at planned start of mating (PSM) and the likelihood of a cow being in calf in first 6 weeks of the mating period³



Thinner cows at PSM have lower pregnancy rates

- Cows that lose 1.0 BCS units after calving were less likely to get in calf in the first six weeks of mating than cows that lose 0.5 BCS unit (**Figure 7**).

Figure 7. Relationship between BCS loss between calving and mating and likelihood of a cow being in calf in first 6 weeks of the mating period ³.



Cows gaining liveweight (Lwt) during the breeding period are more likely to get pregnant, but this effect is small

- Doubling Lwt gain from 0.2 to 0.4 kg/day during breeding only increased 6-week and 12-week in-calf rates by less than 1%³. This equates to lifting BCS gain from 0.25 units/month to 0.5 units/month.

Important points

- Many factors affect reproduction. Body condition is just one of these factors. If your herd's reproductive performance is less than industry targets, use the Reproduction Gap Calculator to assess the value of closing the gaps for your herd.
- Management and/or nutrition have very little effect on BCS loss for the first four to five weeks post-calving.
- Fatter cows at calving cycle earlier but thinner cows lose less BCS after calving. The key is to optimise the benefits of these drivers by calving cows at a BCS of 5.0.
- Younger cows (rising two and three-year-olds) should be 0.5 BCS units fatter than mature cows at calving to optimise reproductive performance.
- To optimise pre-mating cycling and 6-week in-calf rates, cows should calve at a BCS of 5.0 (5.5 for two and three year olds). This will also help limit the BCS loss between calving and mating to no more than 1.0 BCS unit³.

See the InCalf Tools at dairynz.co.nz/support/training/incalf



4. BCS and health

The effects of calving BCS and BCS in early lactation on animal health are inconsistent and depend on the aspect of animal health being considered^{4,5}.

- Metabolic disorders that occur around calving are more prevalent when cows are fat at calving (greater than BCS 6.0), although the risk of milk fever is greater in both fat (greater than BCS 6.0) and thin (less than BCS 3.5) cows.
- The risk of ketosis doubles when calving BCS in mixed aged cows increases from 5.5 to 6.0.
- Calving BCS or change in BCS before calving does not affect the risk of calving difficulty or a stillbirth.
- A difficult calving results in greater BCS loss post-calving.
- Studies have shown that thinner, young cows at calving are more likely to have mastitis and uterine infections in early lactation⁶. The same effect was not evident in mature cows.

There is evidence from more intensive production systems that both low body condition score and loss of body condition score are risk factors for the development of lameness, principally sole and white line disease. While this work has yet to be replicated to the same extent in pasture-based systems, it is likely the same underlying mechanisms would apply in New Zealand conditions. Limiting the rate and extent of body condition score loss may play a role in lameness prevention management on farm⁷.

These data are supportive of BCS targets for first and second calvers being higher than mature cows. To reduce the risk of animal health issues during early lactation, cows should calve at a BCS of 5.0 (5.5 for two- and three-year-olds)^{8,9,10,11}.

5. BCS and other important traits

Body condition score at calving and in early lactation can influence other traits that are important to the profitability and sustainability of the farm system.

- Fatter cows at calving are more likely to give birth to a heifer calf the following year¹². Every unit increase in BCS at calving increases the probability of a heifer calf in the future by approximately 5%.
- Heifers born to cows that lose excessive BCS during early lactation produce less milk and have reduced reproductive capacity⁴.

It is therefore important that a cow is sufficiently fat at calving to maximise replacement heifers, but not so fat as to undermine the productive and reproductive ability of those heifers.

6. BCS and dairy cow welfare

Globally, our understanding of animal welfare has evolved. As well as health and production, we also consider behaviour and affective state, or an animal's experiences and mood. Body condition can be a useful indicator of individual cow welfare^{13,14}, but needs to be considered alongside other factors such as time of year, how quickly she lost condition, and what the farmer is doing to manage the situation.

Generally, the BCS targets that optimise productivity also ensure animal welfare requirements are met. A thin cow may not be in a state of negative welfare provided she is being actively managed but is less robust so stressful situations like transport should be avoided.

The Dairy Cattle Code of Welfare 2019 has identified a minimum acceptable standard of BCS 3.0. The range of BCS within a herd is as important as the average BCS of the herd. When the BCS of any animal falls below 3.0, urgent remedial action must be taken to improve BCS.

7. BCS and profit

A conservative estimate of revenue from increasing cow BCS from 4.0 to 5.0 is \$160 per cow (at \$8.00/kg MS). Improved animal health, greater numbers of replacement heifers the following season, and improvements in the long-term sustainability of the system are not included in this figure.

In comparison, it is estimated that a 500 kg cow requires 125-160 kg of PKE or maize silage to gain a BCS unit while dry (\$45 to \$70 feed expense including wastage).


Factors that affect BCS

1. Time from calving

In pasture-based systems, the profile of BCS change through lactation is generally in the shape of a 'W'. **Figure 8** highlights when genetics and/or management can have an impact on the BCS of a cow during the season.

Figure 8. Typical profile of BCS change in a pasture-based system¹⁵

The black vertical line signifies the point at about 4-5 weeks after calving, when the factors that influence BCS change from wholly genetics to a mix of genetics and management.



The profile of change differs for younger compared to older cows, and for stage of lactation and pregnancy.

First calvers typically calve in better condition than mature cows but take longer to gain the same amount of condition at the end of the season. They need to be dried off earlier than mature cows and need preferential feeding to reach their target calving BCS of 5.5 for their second calving⁴.

Early lactation

Cows are genetically predisposed to lose BCS for the first 50 to 100 days after calving as cow energy intake is less than cow energy requirements. During this period cows are in a natural state of negative energy balance and mobilise body reserves to support maintenance, activity, and milk production. The amount of BCS mobilized during this period is influenced by BCS at calving, cow genetics and, after approximately 4 -5 weeks, management factors.

Mid lactation


Approximately 8-10 weeks after calving, cow dry matter intake peaks, and energy intake matches energy demand. At this stage, cows enter into a positive energy balance and stop mobilising body reserves. This is known as BCS nadir (the lower point of BCS). The rate of change of BCS after nadir depends on genetics and management factors.

Late lactation

After approximately 200 days in milk, cows require energy for pregnancy in addition to maintenance, activity, and milk production. During this period, cows start to partition more energy towards BCS. The amount of BCS gained during this period is influenced by cow genetics and management factors.

Non-lactating (dry) cows

A dry cow uses energy less efficiently than a milking cow for BCS gain, but she does not have the same energy demands for milk production, maintenance, or activity. Therefore, she partitions more energy towards BCS and will gain more BCS per unit of feed eaten (see **page 26** for more on feeding dry cows for BCS gain). The energy demand of pregnancy increases as the cow nears her calving date, such that in the last month before calving, a cow will gain minimal BCS as most of the energy is partitioned to the developing foetus.




2. BCS at calving

Body condition score at calving alters the amount of BCS lost post calving, and therefore influences the BCS that is achieved pre-mating and throughout the season (**Figure 9**).

Note that cows calving at BCS 5.0 tend to return to the same BCS for the next calving. First calvers are an exception to this (see Section 3 - Strategies to achieve targets).

Reaching BCS targets at calving gives your cows the best chance of reaching targets during the season.

Figure 9. Pattern of BCS change for cows calving at different condition scores on research herds in the Waikato¹⁵.



Not all cows will change BCS in the same way when fed the same diet. Cows selected for greater milk production will partition energy towards milksolids at the expense of body condition, resulting in lower BCS throughout lactation^{16,17}.

From these and other studies, we find that:

- Cows that are above BCS targets at calving lose more BCS and are at a greater risk of metabolic diseases such as milk fever and ketosis.
- Cows that are below BCS targets at calving lose less BCS after calving but reach a lower BCS nadir (lowest point) and are at a greater risk of infectious diseases such as metritis and mastitis.
- BCS at calving is the biggest driver of BCS loss after calving and BCS at mating⁴.

3. BCS and genetics

Up to 60% of the variation in BCS within a herd can be due to the genetic makeup of the cows.

During the first 50 to 100 days after calving, BCS loss is largely regulated by genetics. There is very little that can be done from a management perspective to reduce the amount of BCS lost, particularly during the first month after calving.


An emphasis on genetic selection for milk production results in animals that partition more energy to milk production at the expense of BCS.

This has been highlighted in two studies:

The New Zealand Strain Trial¹⁶ compared New Zealand HF cows of 1970s genetic merit (NZ70), with New Zealand HF cows of 1990s genetic merit (NZ90), and HF cows originating from the USA and Europe in the 1990s (NA90). At dry off, NA90 were on average 0.4 BCS units thinner than NZ90 cows, and NZ90 cows were 0.3 BCS units thinner than NZ70 cows.

A later study¹⁷ compared higher producing cows of North American (NA) genetics with lower producing cows of New Zealand (NZ) genetics for the interaction between genetics and feed intake (**Figure 10**). The NA cows continued to lose BCS for longer after calving, reached a lower BCS nadir (the lowest point of BCS), and gained less BCS post-nadir compared to NZ cows, despite addition of supplementary feed.

Figure 10. BCS changes for cows with NZ genetics and North American genetics fed well on pasture only or pasture plus 3 or 6 kg DM supplement/day¹⁷.



Preferential use of energy for milk production, rather than BCS, by different genetic types was highlighted in the same study. NZ cows were able to regain condition during lactation when fed pasture only (**Figure 10a**) and showed a strong BCS response when fed 3 kg DM/day extra feed throughout lactation. By comparison, NA cows struggled to regain condition after reaching their nadir (**Figure 10b**), requiring 6 kg DM/day throughout lactation to put on any significant condition during the rest of lactation.

Recognition of BCS in the BW equation

The BW equation recognises the important role of cow genetics for BCS. The equation penalises animals that naturally lose more than recommended body condition in early lactation and then have to regain it, as these are considered inefficient in a New Zealand system.

4. Nutrition

Body condition score loss during the first four to five weeks post-calving is not primarily influenced by nutrition.

- Cows restricted in intake for the first few weeks post-calving produced less milk, but their loss of BCS and Lwt were not affected^{18, 19}.
- Cows fed high-energy supplements in addition to pasture after calving lost BCS at the same rate as cows that did not receive supplement (**Figure 10**).

Nutrition can influence BCS at four to five weeks after calving.

- Cows fed high energy supplements in addition to pasture reached their BCS nadir earlier and gained more BCS after nadir than cows fed pasture alone (**Figure 10**).
- Genetics still plays a role, such that cows selected for high milk production preferentially partition nutrients to milk production and not to BCS gain, so continue to lose more BCS for a longer period in early lactation¹⁷ (**Figure 10**).

Nutrition continues to influence BCS in mid and late lactation.

- The second trough in the 'W' shape BCS profile curve (**Figure 8**) tends to occur in systems where pasture quality and quantity is limiting. Cows often enter a second period of negative energy balance, usually due to summer dry conditions.
- Cows fed very well on pasture and/or supplemented with concentrates during this period did not experience this second decline in BCS (**Figure 10**).

The lactating cow still preferentially partitions energy towards milk production.

- In the NZ Holstein-Friesian strain trial, Holstein-Friesian cows of NZ origin (NZ90) required 2 kg DM/cow/day maize grain-equivalent for a month (65 kg DM/cow) in addition to pasture, to gain 0.1 BCS units¹⁶. This is equivalent to 1 BCS unit from 650 kg DM maize grain.
- By comparison, Holstein-Friesian cows of North American origin (NA90) required 6.5 kg DM/cow/day maize grain supplement, in addition to pasture, to achieve a similar BCS gain¹⁶. Their response was equivalent to 1 BCS unit from 1,700 kg DM maize grain.

During the dry period, the type of feed offered affects BCS change.

Energy in autumn pasture is used the least efficiently for gaining BCS. Energy from feeds like maize silage, pasture silage and palm kernel extract (PKE) are used 20, 20 and 40% more efficiently, respectively (see **page 28**).

5. Milking frequency

Reducing milking frequency to once a day (OAD) can help increase BCS during lactation. This is because reducing milking to once daily reduces the amount of milk produced, reduces energy demands for walking, and can increase grazing time, thereby providing more energy for condition gain.

In a multi-year farm systems trial²⁰, comparing Jerseys (J) and Holstein-Friesians (HF) that were milked OAD and twice a day (TAD) for whole lactations, cows milked OAD had higher BCS at eight weeks after calving (0.2 to 0.4 BCS units for J and HF, respectively) and at dry off (0.4 to 0.9 BCS units for J and HF respectively). But this was alongside annual milk production losses of 20% for J and 30% for HF.

Strategic use of OAD milking in early lactation can alter the energy status of cows after calving^{21,22}. A short period of OAD milking after calving led to OAD milked cows being slightly higher in BCS (0.15 to 0.23 BCS units for three and six weeks of OAD milking, respectively) than cows milked TAD, by six weeks after calving, and they remained higher through mating and mid lactation. Production losses were in the order of 15-20% during OAD milking, and remained lower when cows returned to TAD, leading to seven and 12% less total milksolids production by late lactation. When there is a feed shortage in early spring, OAD milking after calving can improve energy status of cows receiving less than their ideal ration, but milk production will be impacted²³.

Milking cows three times in two days (3in2) has less impact on milk production, with only a 5% reduction in full season production compared to TAD milking. Whole season farmlet studies of 3in2 milking found the energy saved from walking was sufficient for a 0.25 unit BCS gain over the whole lactation for cows milked 3in2 compared to TAD. Use on commercial farms has revealed lower production losses and greater BCS gains can be achieved.

For more information, go to dairynz.co.nz/milking/milking-intervals

See **Section 3** - Strategies to achieve targets for more information on how to use milking frequency in late lactation to improve BCS.



Summary

Remember that BCS targets at calving have been set to manage the balancing act between optimising dry matter intake, milk production and reproduction, and minimising health and welfare issues of being too fat, or too thin. Achieving BCS targets at calving can have far-reaching impacts for individual cows, and your herd.

Cows that are below target BCS at calving tend to:

- produce less milksolids during lactation,
- cycle later after calving,
- be more prone to milk fever, mastitis, uterine infections, and lameness, and
- are more likely to become pregnant with bull calves, at their next mating.

Cows that are above target BCS at calving tend to:

- eat less after calving,
- be more prone to milk fever, ketosis, and metabolic diseases after calving,
- and lose more condition after calving.

This in turn can affect:

- their ability to get in calf at mating time, and
- the lifetime production and reproductive success of their daughters.

The change in BCS of individual cows through lactation is affected by:

- Time since calving,
- BCS at calving,
- Genetic makeup of the cow,
- Type of feed and amount of feed, and
- Milking frequency.

References

Section 3

1. Grala, T. M., R. R. Handley, J. R. Roche, C. G. Walker, C. V. C. Phyn, and J. K. Kay. 2016. Once-daily milking during late lactation in pasture-fed dairy cows has minor effects on feed intake, body condition score gain, and hepatic gene expression. *Journal of Dairy Science* 99:3041-3055.
2. Edwards, J.P., N. McMillan, R. Bryant, and B. Kuhn-Sherlock. 2020. Milking options to increase flexibility and workplace attractiveness. DairyNZ project summary report. Accessed May 2021 from: dairynz.co.nz/media/5794325/milking_options_to_increase_flexibility_and_workplace_attractiveness_v2.pdf.
3. Mandok, K. M., J. K. Kay, S. L. Greenwood, J. P. McNamara, M. Crookenden, R. White, S. Shields, G. R. Edwards, and J. R. Roche. 2014. Efficiency of use of metabolizable energy for body weight gain in pasture-based, nonlactating dairy cows. *Journal of Dairy Science* 97:4639-4648.
4. Kay, J. Feeding for condition score gain. 2014. DairyNZ Technical Series, April, Issue 21: 5-8. Accessed May 2021 from: dairynz.co.nz/media/424995/technical_series_april_2014.pdf
5. Mandok, K. S., J. K. Kay, S. L. Greenwood, G. R. Edwards, and J. R. Roche. 2013. Requirements for zero energy balance of nonlactating, pregnant dairy cows fed fresh autumn pasture are greater than currently estimated. *Journal of Dairy Science* 96:4070-4076.
6. Edwards, G. R., J. M. de Ruiter, D. E. Dalley, J. B. Pinxterhuis, K. C. Cameron, R. H. Bryant, H. J. Di, B. J. Malcolm, and D. F. Chapman. 2014. Dry matter intake and body condition score change of dairy cows grazing fodder beet, kale and kale-oat forage systems in winter. *Proceedings of the New Zealand Grassland Association* pp 81-88

Section 4

1. Roche, J. R., D. Blache, J. K. Kay, D. R. Miller, A. J. Sheahan, and D. W. Miller. 2008. Neuroendocrine and physiological regulation of intake with particular reference to domesticated ruminant animals. *Nutrition Research Reviews* 21:207-234.
2. Roche, J. R., J. M. Lee, K. A. Macdonald, and D. P. Berry. 2007. Relationships among body condition score, body weight, and milk production variables in pasture-based dairy cows. *Journal of Dairy Science* 90:3802-3815.
3. Roche, J. R., K. A. Macdonald, C. R. Burke, J. M. Lee, and D. P. Berry. 2007. Associations among body condition score, body weight, and reproductive performance in seasonal-calving dairy cattle. *Journal of Dairy Science* 90:376-391.
4. Roche, J. R., N. C. Friggens, J. K. Kay, M. W. Fisher, K. J. Stafford, and D. P. Berry. 2009. Invited review: Body condition score and its association with dairy cow productivity, health, and welfare. *Journal of Dairy Science* 92:5769-5801.
5. Roche, J. R., K. A. Macdonald, K. E. Schutz, L. R. Matthews, G. A. Verkerk, S. Meier, J. J. Loor, A. R. Rogers, J. McGowan, S. R. Morgan, S. Taukiri, and J. R. Webster. 2013. Calving body condition score affects indicators of health in grazing dairy cows. *Journal of Dairy Science* 96:5811-5825.
6. McDougall, S., H. Hussein, D. Aberdein, K. Buckle, J. Roche, C. Burke, M. Mitchell, and S. Meier. 2011. Relationships between cytology, bacteriology and vaginal discharge scores and reproductive performance in dairy cattle. *Theriogenology* 76:229-240.
7. Randall, L. V., M. J. Green, L. E. Green, M. G. G. Chagunda, C. Mason, S. C. Archer, and J. N. Huxley. 2018. The contribution of previous lameness events and body condition score to the occurrence of lameness in dairy herds: A study of 2 herds. *Journal of Dairy Science* 101:1311-1324.
8. Crookenden, M. A., C. G. Walker, A. Heiser, A. Murray, V. S. R. Dukkipati, J. K. Kay, S. Meier, K. M. Moyes, M. D. Mitchell, J. J. Loor, and J. R. Roche. 2017. Effects of precalving body condition and prepartum feeding level on gene expression in circulating neutrophils. *Journal of Dairy Science* 100:2310-2322.
9. Lange, J., A. McCarthy, J. Kay, S. Meier, C. Walker, M. A. Crookenden, M. D. Mitchell, J. J. Loor, J. R. Roche, and A. Heiser. 2016. Prepartum feeding level and body condition score affect immunological performance in grazing dairy cows during the transition period. *Journal of Dairy Science* 99:2329-2338.
10. Roche, J., A. Bell, T. Overton, and J. J. Loor. 2013. Nutritional management of the transition cow in the 21st century—a paradigm shift in thinking. *Animal Production Science* 53:1000-1023.

11. Roche, J. R., S. Meier, A. Heiser, M. D. Mitchell, C. G. Walker, M. A. Crookenden, M. V. Riboni, J. J. Looor, and J. K. Kay. 2015. Effects of precalving body condition score and prepartum feeding level on production, reproduction, and health parameters in pasture-based transition dairy cows. *Journal of Dairy Science* 98:7164-7182.
12. Roche, J. R., J. M. Lee, and D. P. Berry. 2006. Pre-conception energy balance and secondary sex ratio - partial support for the Trivers-Willard hypothesis in dairy cows. *Journal of Dairy Science* 89:2119-2125.
13. Matthews, L. R., C. Cameron, A. J. Sheahan, E. S. Kolver, and J. R. Roche. 2012. Associations among dairy cow body condition and welfare-associated behavioral traits. *Journal of Dairy Science* 95:2595-2601.
14. Schutz, K. E., N. R. Cox, K. A. Macdonald, J. R. Roche, G. A. Verkerk, A. R. Rogers, C. B. Tucker, L. R. Matthews, S. Meier, and J. R. Webster. 2013. Behavioral and physiological effects of a short-term feed restriction in lactating dairy cattle with different body condition scores at calving. *Journal of Dairy Science* 96:4465-4476.
15. Roche, J. R., D. P. Berry, J. M. Lee, K. A. Macdonald, and R. C. Boston. 2007. Describing the body condition score change between successive calvings: a novel strategy generalizable to diverse cohorts. *Journal of Dairy Science* 90:4378- 4396
16. Macdonald, K. A., G. A. Verkerk, B. S. Thorrold, J. E. Pryce, J. W. Penno, L. R. McNaughton, L. J. Burton, J. A. Lancaster, J.H. Williamson, and C. W. Holmes. 2008. A comparison of three strains of Holstein-Friesian grazed on pasture and managed under different feed allowances. *Journal of Dairy Science* 91:1693-1707.
17. Roche, J. R., D. P. Berry, and E. S. Kolver. 2006. Holstein-Friesian strain and feed effects on milk production, body weight, and body condition score profiles in grazing dairy cows. *Journal of Dairy Science* 89:3532-3543.
18. Roche, J. R. 2007. Milk production responses to pre- and postcalving dry matter intake in grazing dairy cows. *Livestock Science* 110:12-24.
19. Roche, J., C. Burke, S. Meier, and C. Walker. 2011 Reproductive failure – do cows need more feed to get in calf? DairyNZ Technical Series, September, Issue 5:2-5. Accessed May 2021 at: https://www.dairynz.co.nz/media/424963/technical_series_september_2011.pdf
20. Clark, D., C. Phyn, M. Tong, S. Collis, and D. Dalley. 2006. A systems comparison of once-versus twice-daily milking of pastured dairy cows. *Journal of Dairy Science* 89:1854-1862.
21. Phyn, C., J. K. Kay, A. Rius, T. Grala, and J. Roche. 2013 Early lactation milking affects season's production. DairyNZ Technical Series, June, Issue 16:4-8. Accessed May 2021 at: dairynz.co.nz/publications/technical-series/technical-series-june-2013
22. Phyn, C. V. C., J. K. Kay, A. G. Rius, S. R. Morgan, C. G. Roach, T. M. Grala, and J. R. Roche. 2014. Temporary alterations to post- partum milking frequency affect whole-lactation milk production and the energy status of pasture- grazed dairy cows. *Journal of Dairy Science*. 97:6850–6868.
23. Kay, J. K., C. V. C. Phyn, A. G. Rius, S. R. Morgan, T. M. Grala, and J. R. Roche. 2013. Once-daily milking during a feed deficit decreases milk production but improves energy status in early lactating grazing dairy cows. *Journal of Dairy Science*. 96:6274–6284.

DairyNZ 

dairynz.co.nz